




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Journal Pre-proof

Re: 'High prevalence of heterotopic ossification in critically ill patients with severe COVID-19' by Stoira et al

Nicolas de l'Escalopier, Laurent Mathieu, Christophe Duret, Sébastien Banzet, François Genêt, Marjorie Salga



PII: S1198-743X(21)00030-6

DOI: <https://doi.org/10.1016/j.cmi.2020.12.038>

Reference: CMI 2390

To appear in: *Clinical Microbiology and Infection*

Received Date: 26 November 2020

Accepted Date: 1 December 2020

Please cite this article as: de l'Escalopier N, Mathieu L, Duret C, Banzet S, Genêt F, Salga M, Re: 'High prevalence of heterotopic ossification in critically ill patients with severe COVID-19' by Stoira et al, *Clinical Microbiology and Infection*, <https://doi.org/10.1016/j.cmi.2020.12.038>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 European Society of Clinical Microbiology and Infectious Diseases. Published by Elsevier Ltd. All rights reserved.

Re: 'High prevalence of heterotopic ossification in critically ill patients with severe COVID-19' by Stoira et al.

Nicolas de l'Escalopier^{1,2}, Laurent Mathieu^{1,3}, Christophe Duret², Sébastien Banzet^{4,5}, François Genêt^{6,7} and Marjorie Salga^{6,7}

1. Department of orthopedic, trauma and reconstructive surgery, Percy Military Hospital, 101 avenue Henri Barbusse, 92140 Clamart, France
2. CRF Les Trois Soleils, Médecine Physique et de Réadaptation, Unité de Rééducation Neurologique, Boissise-Le-Roi (77) France
3. Department of surgery, French Military Health Service Academy, Ecole du Val-de-Grâce, Paris, France
4. French Military Institute for Biomedical Research, Cell and Tissue Therapy unit, Clamart, France
5. Inserm UMR-MD-1197, Clamart, France
6. Department of Physical Medicine and Rehabilitation, CIC 1429, Raymond Poincaré Hospital, APHP, Garches, France.
7. END:ICAP U1179 INSERM, University of Versailles Saint Quentin en Yvelines, UFR Simone Veil-Santé, Montigny le Bretonneux, France.

*Corresponding author ndelescalopier@gmail.com

Dear editor,

We would like to respond to the letter to the editor sent by Stoiria et al. « High prevalence of heterotopic ossification in critically ill patients with severe COVID-19 » to give the point of view of an experienced medico-surgical team in charge of neurogenic Heterotopic Ossification (HO) case management [1].

Like Stoiria et al., we have experienced a high prevalence of troublesome HO in severe COVID patients. Another rehabilitation team described HO located around the same joints after COVID [2]. Meanwhile in our service, already 3 surgeries have been performed to remove HO located around the hip. Two of them were anterior to the hip, while the third was posterior and responsible for sciatic nerve compression. Additionally, we have removed another HO of the inner elbow that was in contact with the ulnar nerve. Interestingly, development of HO in patients treated for SARS-CoV in 2003 has been reported. We believe that HO formation is likely due to the severe and prolonged inflammation experienced during intensive care, perhaps more than the virus itself [3]. Future observations of dexamethasone use to manage severe COVID patients will provide information about pathophysiological inflammatory mechanisms. HO occurs more frequently in patients who develop intensive care unit-acquired neuromuscular abnormalities, and the role of peripheral neurologic system damage has already been described for neurogenic HO [4]. Stoiria et al assume that prolonged immobilization is key in HO occurrence, congruent with the observation that curare use is a risk factor for HO after severe burns [5]. Preventive physiotherapy measures have been described, including daily joint mobilization as early as possible.

They are many similarities between HO after COVID and neurogenic HO: developmental, morphological, symptoms and risks. Therefore, there is no reason not to manage COVID HO like classical neurogenic HO. The important points to consider are well defined [1]. Determining if HO is troublesome is the first step before treatment, and non-disabled patients that potentially can recover full autonomy may be more sensitive to HO sequelae. If troublesome, pain cannot be alleviated non-surgically, or signs of neural or vascular compression are present, surgery becomes necessary. Early surgery is feasible and should be performed to avoid complications. However, resection does not need to be complete as only the problematic zone of the HO mass needs to be removed. Contrast-enhanced CT with 3-dimensional reconstruction provides critical mapping of coinciding blood vessels and nerves, which is critical for determining a surgical strategy.

The report by Stoiria *et al.*, establishes that HO in severe COVID patients develop more often than expected. Survivors of severe COVID that develop HO may experience complications as a result. These patients should be referred to a medico-surgical team specialized in HO for treatment, as strategies to manage HO are well established.

- [1] Denormandie P, de l'Escalopier N, Gatin L, Grelier A, Genêt F. Resection of neurogenic heterotopic ossification (NHO) of the hip. *Orthop Traumatol Surg Res* 2017.
- [2] Meyer C, Haustrate M-A, Nisolle JF, Deltombe T. Heterotopic ossification in COVID-19: A series of 4 cases. *Ann Phys Rehabil Med* 2020.
- [3] Genêt F, Kulina I, Vaquette C, Torossian F, Millard S, Pettit AR, et al. Neurological heterotopic ossification following spinal cord injury is triggered by macrophage-mediated inflammation in muscle. *J Pathol* 2015;236:229–40.
- [4] Debaud C, Salga M, Begot L, Holy X, Chedik M, de l'Escalopier N, et al. Peripheral denervation participates in heterotopic ossification in a spinal cord injury model. *PLoS One* 2017;12:e0182454.
- [5] Thefenne L, de Brier G, Leclerc T, Jourdan C, Nicolas C, Truffaut S, et al. Two new risk factors for heterotopic ossification development after severe burns. *PLoS One* 2017;12:e0182303.